

Begonia semperflorens F₁

Sprint Plus

White

Art.-Nr.: BS0206P



- Traditional compact semperflorens
- Very early flowering
- Perfectly suitable in pots and packs
- Fast and uniform growing
- Green foliage partner to Nightlife

Crop Time	Frühjahr: 9 - 11 Wochen
Höhe 	23 cm
Breite 	28 cm
Standort	Sun - Shade
Seed Form	Pilliert
Product Use	Topfpflanze, Ampel, Mixed Containers, Landschaft
Family, Origin	Begoniaceae, South and Central America
Minimum Germ. Rate	90 %

TECHNICAL GUIDE

Begonia semperflorens F₁ Sprint Plus

Verwendung

Packs, Pots, Hanging Baskets, Mixed Containers and Landscape

Kornablage

1-2 seeds or pellets per plug

Keimbedingung

Optimum conditions for seedling development, beginning on the day of sowing until radicle emergence. Expect radicle emergence in 6-8 days.

Medien

Plug Culture: pH 5.5-5.8; EC 0.5-0.75.

Growing On: pH 5.5-5.8; EC 1.25-1.5.

Temperatur

Plug Culture: 72–76°F (22–24°C) days 1–11. For irrigation use warm water (above 64°F / 18°C) only.

Growing on: 68–70°F (20–21°C) nights, 64–67°F (18–19°C) days for the first 14 days or until the roots reach the bottom of the container. Thereafter temperatures may be lowered to 62–65°F (16–18°C) day and night. An ADT (average daily temperature of 67°F (19°C) will give the fastest finished crop.

Düngung

Plug Culture: Maintain an EC < 1.0. Fertilized water should not exceed an EC of 0.5. Initial feeding should be with a balanced fertilizer low in ammonium. Begin feeding with a 14-4-14, 14-2-14 or 17-5-17 fertilizer at 50–60 ppm.

Growing On: Moderate fertilization levels are required. Fertilize the crop weekly with 100-150 ppm nitrogen, using a complete balanced fertilizer. Avoid high ammonium and high nitrogen levels, because the foliage can grow very large. Avoid pH levels above 6.0, as high pH can cause iron deficiency. Watch for low Ca and Mg levels since this can result in stunted plants with marginal leaf edge burn. Under high light conditions use an ammonium based fertilizer (17-5-17) and under low light use a calcium based fertilizer (14-4-14).

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

FARBEN DER SERIE

Begonia semperflorens F₁ Sprint Plus



White
BS0206P



Orange
BS0201P



Orange Bicolor
BS0202P



Lipstick
BS0204P



Red
BS0203P



Pink
BS0207P



Rose
BS0205P



Blush
BS0208P



Maxi Mix
BS0299P