

Celosia cristata

Brainiac

Robo Red

Art.-Nr.: CC4005R



- Big, dense and early flowering heads
- Compact and uniform plant habit
- Tolerates hot conditions in moderate climates
- Recommended for pots and packs

Crop Time	Frühjahr: 8 - 10 Wochen
Höhe 	15 cm
Breite 	18 cm
Standort	Sun
Seed Form	Roh
Product Use	Packs, Topfpflanze, Mixed Containers, Landschaft
Minimum Germ. Rate	85%

TECHNICAL GUIDE

Celosia cristata Brainiac

Verwendung

Mass plantings, specimen plant and mixed containers

Aussaattermin

January-June

Kornablage

1 seed per plug

Keimbedingung

Stage I: 7-10 days at 25 °C; Stage II-IV: 11-20 days at 18-21 °C day temperature and 15-18 °C night temperature. Use media with low soluble salt levels and pH: 5.5-6.0. Requires light for germination. Cover seed lightly with vermiculite after sowing. Keep soil slightly moist but not wet.

Medien

Use a well-drained, growing substrate with 15-30 % clay, 1-1,5 kg/m³ complete balanced fertilizer, iron-chelate, micronutrients, pH: 5.5-6.0.

Temperatur

Grow at 18-20 °C. At the end of the cultivation time decrease the temperature to 16 °C. Avoid temperatures below 16 °C, because Celosia needs warm temperatures to grow.

Düngung

Moderate fertilization levels are required. Fertilize the crop weekly with 150-200 ppm nitrogen, using a potassium and phosphate balanced fertilizer (N: K₂O-Ratio: 1 :1,5). Avoid high ammonium and high nitrogen levels. Prevent magnesium deficiency by applying magnesium sulphate (0,05 %) 1-2 times and in case of iron deficiency apply iron-chelate for 1-2 times.

Expert Tip

Cultivate plants firstly slightly moist and later dry. Avoid an inhibited and stressful cultivation (in particular low temperatures or nutrition deficiency), because the flower initiation may start before plants have reached sufficient size and thus the plant quality decreases.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

FARBEN DER SERIE

Celosia cristata Brainiac



Lightning Yellow
CC4001R



Mad Magenta
CC4002R



Raven Red
CC4003R



Robo Red
CC4005R



Think Pink
CC4004R



Mix (excl. Raven Red)
CC4099R