

Delphinium grandiflorum

Blue Mirror

Art.-Nr.: DG0101R



- FastraX perennial: First year flowering without vernalization
- Intense blue color
- Full, round plant habit
- Perfect for beds and borders

Crop Time	Frühjahr: 16 - 20 Wochen
Höhe ♂	50 cm
Standort	Sun - Partial shade
Seed Form	Roh
Heat Zone	6-1
Hardiness Zone	3b-7b
Product Use	Beet und Balkon, Landschaft

TECHNICAL GUIDE

Delphinium grandiflorum Blue Mirror

Verwendung

Bedding and border plants, pot and container plants

Aussaattermin

Annual forcing: December-May; Perennial forcing: Mid August-Mid September including wintering

Kornablage

2-3 seeds per plug

Keimbedingung

14-18 days at temperatures of 68-72 °F (20-22 °C), cover seed lightly and maintain constant humidity levels. Stage II: gradually lower temperatures to 64-68 °F (17-20 °C). Stage III & IV: begin light fertilization at 50-75 ppm nitrate nitrogen. Use low rates of phosphorus during plug production

Topfen

Transplant plugs 6-8 weeks after sowing. Finish at 45-75 °F (7-24 °C)

Medien

Use a well-drained, growing substrate with 10-30 % clay, 0-15 % parts (e.g. bark, sand perlite) 1-2 kg/m³ complete balanced fertilizer, 2-4 kg/m³ slow release fertilizer (3-9 months), iron-chelate, micronutrients, pH: 5.8-6.5.

Temperatur

Grow at 13-16 °C or outdoors. In winter indoors frost free at 3-5 °C or outdoors. Outdoor fleece cover and dry substrate needed. In spring the plants start to grow for 8-10 weeks at 15-20 °C. Temperatures below 13 °C in combination with short day is a cause for rosette leaf formation and inhibition of the flower initiation.

Düngung

Moderate fertilization levels are required. Fertilize the crop weekly with 80-100 ppm nitrogen (at 2 kg/m³ slow release fertilizer in substrate), using a complete balanced fertilizer. Avoid high ammonium and high nitrogen levels. Don't fertilize after flowering or after mid September. In spring fertilize with 130-150 ppm nitrogen using a complete balanced fertilizer. Prevent magnesium deficiency by applying magnesium sulphate (0,05 %) 1-2 times and in case of iron deficiency apply iron-chelate for 1-2 times. The fertilization with slow release fertilizer is favourable as against with complete balanced fertilizer, because the roots are very sensitive to high salt levels in substrates. Avoid high fertilizer concentrations, it is advisable to fertilize several times with low concentrations weekly.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

FARBEN DER SERIE

Delphinium grandiflorum Blue Mirror



DG0101R