

Helianthus annuus

# Pacino®

**Gold**

Art.-Nr.: HA1101R



- Compact, pollenless pot sunflower
- Different, vibrant yellow color tones
- High quantity of small flowers

**Technical Guide:** [Click here](#)

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

<b>Crop Time</b>	Frühjahr: 8 - 10 Wochen
<b>Höhe</b> ∅	33 cm
<b>Breite</b> ∅	23 cm
<b>Flower Size</b> ∅	11 cm
<b>Standort</b>	Sun
<b>Seed Form</b>	Roh
<b>Best Uses</b>	Beet und Balkon, Topfpflanze

## KULTURANLEITUNG

Helianthus annuus Pacino®

### Verwendung

Plants for bedding, attractive pot plants and container plants for patio.

### Aussaattermin

March-End July, sowing in intervals

### Kornablage

1-2 seeds per plug, can be sown directly into final pot

### Keimbedingung

5 days at 65-75 °F (18-24 °C). Stage II 70 °F (21 °C). Supplemental lighting during germination and plug production to control seedling stretch is beneficial to overall crop quality. For Stage III & IV reduce temperatures to 62 °F (17 °C), and begin feeding at 100-150 ppm nitrogen.

### Topfen

Direct sow seed or transplant at two true leaf stage. Continue feeding at 100 ppm nitrogen weekly. Maintain temperatures above 62 °F (17 °C) for improved branching and overall quality. Fertilize weekly at 200-250 ppm nitrogen. Avoid excessive phosphorus.

### Medien

Use a well-drained, growing substrate with 15-30 % clay, 1-3 kg/m<sup>3</sup> complete balanced fertilizer, 0-2 kg/m<sup>3</sup> slow release fertilizer (3-6 months), iron-chelate, micronutrients, pH: 5.5-6.2.

### Temperatur

Grow at 15-18 °C. Cultivate flowering plants not below 12 °C. Helianthus plants do not tolerate frost.

### Düngung

High fertilization levels are required. Fertilize the crop weekly with 200-250 ppm nitrogen, using a potassium balanced fertilizer (N: K<sub>2</sub>O-ratio: 1:1,5). Avoid high ammonium and high nitrogen levels. Prevent magnesium deficiency by applying magnesium sulphate (0,05 %) 1-2 times and in case of iron deficiency apply iron-chelate for 1-2 times.

---

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

## FARBEN DER SERIE

Helianthus annuus Pacino<sup>®</sup>



**Cola**  
HA1102R



**Gold**  
HA1101R



**Lemon**  
N0850



**Mix (incl. Lemon)**  
HA1199R