Lavandula angustifolia

Munstead



- Deep lavender blue flowers
- Compact evergreen clump-forming perennial

Crop Time	Frühjahr: 23 - 25 Wochen
Höhe ⌀	40 cm
Standort	Sun
Seed Form	Roh, ApeX
Heat Zone	12-7
Hardiness Zone	5a-9a
Product Use	Beet und Balkon, Landschaft

TECHNICAL GUIDE

Lavandula angustifolia Munstead

Verwendung
Plants for border and open positions, ornamental leaf plant, aromatic plant, medicinal plant, pot and container plants, plants attractive for bees, extensive roof planting

Aussaattermin
January-March for green pots, June-August for flowering in pots the following year

Kornablage
Optimal storage ApeX: Up to 6 months at 5 °C / 41 °F

1-2 seeds per plug

Keimbedingung
Germinates in 14-20 days at 65-72 °F (18-22 °C). Light is required for germination. Treated ApeX seed increases germination percentages over that of raw seed.

Topfen
Transplant plugs after 9 weeks. Grow on at 55-65 °F (12-18 °C) in an alkaline medium. Vernalization is required for flower initiation. After vernalization, begin forcing plants at temperatures of 60-65 °F (15-18 °C) for 6-7 weeks under long days to promote flowering.

Media
Use a well-drained, growing perennial substrate with 0-15 % clay, 0-15 % parts (e.g. perlite, sand, bark), 1-2 kg/m^3 complete balanced fertilizer, 0-2 kg/m^3 slow release fertilizer (3-9 months), iron-chelate, micronutrients, pH: 5.8-7.0.

Temperatur
Grow at 12-21 °C or outdoors. In winter indoors frost free at 3-5 °C or outdoors. Outdoor fleece cover needed. In spring the plants start to grow for 6-8 weeks at 18-21 °C. Cold temperatures at 12-15 °C will increase the cultivation time by 3-4 weeks. A chilling period (vernalization) for flower initiation for at least 15 weeks at 3-5 °C is required. A forcing in spring with warm temperatures need high light intensity and decreases the cultivation time.

Düngung
Moderate fertilization levels are required. Fertilize the crop weekly with 100-150 ppm nitrogen (at 2 kg/m^3 slow release fertilizer in substrate), using a complete balanced fertilizer. Avoid high ammonium and high nitrogen levels. High nitrogen levels are the cause that the shoots are stretched and the plant fall apart. Don't fertilize after mid September. In spring fertilize 100-150 ppm nitrogen of a complete balanced fertilizer. Prevent magnesium deficiency by applying magnesium sulphate (0,05 %) 1-2 times and in case of iron deficiency (above pH 6.0) apply iron-chelate for 1-2 times.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.
Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.
Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.
Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

FARBEN DER SERIE

Lavandula angustifolia Munstead



Variety
LA0303R/T