

Lobelia speciosa F<sub>1</sub>

**Fan**®

**Burgundy**

Art.-Nr.: LS2002P



- FastraX perennial: First year flowering without vernalization
- Excellent branching of uniform plants
- Attractive red-green leaves of Scarlet & Scarlet Compact
- Fantastic germination: 90%

<b>Crop Time</b>	Frühjahr: 18 - 20 Wochen
<b>Höhe</b> ☞	60 cm
<b>Breite</b> ☞	35 cm
<b>Standort</b>	Sun - Partial shade
<b>Seed Form</b>	Pilliert
<b>Heat Zone</b>	9-2
<b>Hardiness Zone</b>	6-10
<b>Product Use</b>	Topfpflanze, Containers, Beet und Balkon
<b>Family, Origin</b>	Campanulaceae, (sub)tropical region
<b>Minimum Germ. Rate</b>	85%

## TECHNICAL GUIDE

Lobelia speciosa F<sub>1</sub> Fan®

### Verwendung

Pots, Containers and Beds

### Aussaattermin

January-March for flowering in pots from July onwards, August-October for flowering in pots the following year, April-May for cut flower production outdoors

### Kornablage

1 seed per plug for short day conditions 3-5 seeds per plug for long day conditions, because in long day the plants do not form a leaf rosette.

### Keimbedingung

Optimum conditions for seedling development, beginning on the day of sowing until radicle emergence. Expect radicle emergence in 8-12 days.

### Topfen

Transplant plugs after 10 weeks. Transplant one plant into a 4-6" (10-15 cm) pot, or 2-3 plants in a 6" (15 cm) or larger container. Apply a well balanced fertilizer every 2 weeks. Grow on at 60-65 °F (15-18 °C). Vernalization is not required for flower initiation.

### Medien

Plug culture: pH 5.8-6.2. EC 0.7-1.2.

Growing on: pH 5.8-6.4. EC 1.1-1.3.

### Temperatur

Plug culture: 20-22 °C (68-72 °F) until radicle emergence. Afterwards, ensure 18-20 °C (64-68 °F) during night and day. When the roots reach the bottom of the cell, the temperature can be lowered to 16-18 °C ( 60-64 °F).

Growing on: Maintain 16-18 °C ( 60-64 °F) during night and day. Once established in the final container, the temperature can be lowered to 10-15 °C (50-59 °F). Cooler temperatures support the uniformity, stability and compactness of the plants, even if the crop time increases somewhat. An outdoor production is also possible. In winter, a frost-free cultivation indoors at 3-5 °C (38-40 °F) or an outdoor cultivation with a fleece cover is possible. The plants are sensitive to strong frost temperatures.

### Düngung

Moderate-high fertilization levels are required. Fertilize the crop weekly with 150-200 ppm nitrogen (at 0 kg/m<sup>3</sup> slow release fertilizer in substrate), using a complete balanced fertilizer. Avoid high ammonium and high nitrogen levels. Don't fertilize after mid September. In spring fertilize with 100-150 ppm nitrogen of a complete balanced fertilizer. Prevent magnesium deficiency by applying magnesium sulphate (0,05 %) 1-2 times and in case of iron deficiency apply iron-chelate for 1-2 times. Field: If necessary according to analysis, improve the soil with 80-100 g/m<sup>2</sup> of a slow release fertilizer per year, applied in several portions.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

## FARBEN DER SERIE

Lobelia speciosa F<sub>1</sub> Fan®



**Burgundy**  
LS2002P



**Scarlet**  
LS2001P



**Scarlet Compact**  
LS2003P