

Lobelia erinus

Palace

Mix

Art.-Nr.: LE0199R/U



- Early flowering series
- Compact plant habit
- Excellent for pots, baskets and borders

In North America only Multipellets available.

Crop Time	Frühjahr: 9 - 11 Wochen
Höhe 	18 cm
Breite 	23 cm
Standort	Partial shade
Seed Form	Roh, Multipille
Product Use	Beet und Balkon, Ampel, Topfpflanze

TECHNICAL GUIDE

Lobelia erinus Palace

Verwendung

Beds, borders, containers and color bowls

Kornablage

7-10 seeds or 1 multipellet per plug (multi-seed pellets produce an average of 5-7 seedlings per pellet)

Keimbedingung

Stage I: 4-7 days at 70-75 °F (21-24 °C), in media with very low soluble salt levels and pH 5.5-6.5. Requires light for germination. Do not cover seeds. If pellets are used, ensure that the pellets are well-watered to allow them to dissolve. Keep soil slightly moist but not wet. Long day (12-14 h) during germination decreases the cultivation time and flowering will be earlier. Avoid direct sunlight by shading seeds after sowing.

Topfen

4-5 weeks after sowing, transplant into packs or pot 6-10 cm (2,5-4").

Medien

Use a well-drained, disease-free, soilless medium with a pH 5.2 -5.8.

Temperatur

Grow at night temperatures of 12-15 °C (54-59 °F) and day 17 -20 °C (63-68 °F). After development of the roots temperature can be further decreased. Before selling harden the plants by decreasing the temperature to 6-8 °C (43-46 °F). Lobelia erinus does not tolerate frost.

Düngung

Moderate fertilization levels are required. Fertilize the crop weekly with 150-200 ppm nitrogen, using a complete balanced fertilizer. Avoid high ammonium and high nitrogen levels. Wetness and cold medium temperatures are a cause for iron deficiency. The roots are sensitive to high salt levels in substrates. Avoid high fertilizer concentrations, it is advisable to fertilize several times with low concentrations weekly. Reduce fertilizer rate at the end of the crop to keep the plants compact.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

FARBEN DER SERIE

Lobelia erinus Palace



White
LE0104R/U



Blue
LE0101R/U



Blue with Eye
LE0102R/U



Lilac
LE0103R/U



Royal
LE0106R/U



Sky Blue
LE0105R/U



Mix
LE0199R/U