

Portulaca grandiflora F₁

Stopwatch

Cream

Art.-Nr.: PG2001R/U



- Compact, mounding habit - plants grow more upright
- Early flowering
- Tighter habit for packs

Technical Guide: [Click here](#)

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

Crop Time	Frühjahr: 10 - 12 Wochen
Höhe 	12 cm
Breite 	28 cm
Standort	Sun
Seed Form	Roh, Multipille
Best Uses	Beet und Balkon, Landschaft, Steingarten

KULTURANLEITUNG

Portulaca grandiflora F₁ Stopwatch

Verwendung

Mixed containers, bedding, landscaping

Kornablage

2 seeds per plug, 1 multi-seed pellet per plug

Keimbedingung

Stage I: 2-3 days at 70-78 °F (22-26 °C). Sow in media with very low soluble salt levels and pH 5.5-6.2. Light is not essential, but beneficial for germination. Keep soil moist but not wet. Keep relative humidity between 90-95%. At the end of stage I the radicle will have emerged and the cotyledons begin to unfurl.

Topfen

4-5 weeks after sowing (288-tray) transplant 1 plant into 10-11 cm (4-5") pots or packs

Medien

Use a well-drained substrate with pH 5.5 -6.2 and 1-1.5kg/m³ fertilizer

Temperatur

18 °C-22 °C (65-72 °F). Temperatures below 18 °C (65 °F) will inhibit induction and result in poor or no vegetative growth.

Düngung

Moderate fertilization levels are required. Fertilize the crop with 150-200 ppm nitrogen, using a complete balanced fertilizer.

Tip

Our Portulaca series gets our growers more turns since it is at least 10 days earlier to flower with a well branched habit that fills pots & packs quickly.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

FARBEN DER SERIE

Portulaca grandiflora F₁ Stopwatch



Cream
PG2001R/U



Fuchsia
PG2002R/U



Orange
PG2003R/U



Rose
PG2004R/U



Scarlet
PG2005R/U



White
PG2006R/U



Yellow
PG2007R/U



Mix
PG2099R/U