


Delphinium hybrida

# Benarys Pacific

**Cameliard**

Art.-Nr.: DH0102R



- FastraX perennial: First year flowering without vernalization
- Rigorously tested for excellent flower and stem quality
- Excellent cut flower with long, impressive flower stems
- Long vase life

<b>Crop Time</b>	Frühjahr: 18 - 22 Wochen
<b>Höhe</b> 	160 cm
<b>Standort</b>	Sun
<b>Seed Form</b>	Roh
<b>Heat Zone</b>	6-1
<b>Hardiness Zone</b>	3b-7b
<b>Best Uses</b>	Beet und Balkon, Schnittblume, Landschaft

## KULTURANLEITUNG

Delphinium hybrida Benarys Pacific

### Verwendung

Background plant, cut flower

### Aussaattermin

Outdoor forcing: March-April; Indoor forcing: December-May

### Kornablage

2-3 seeds per plug

### Keimbedingung

20-30 days at temperatures of 59-68 °F (15-18 °C).

### Topfen

Grow on at 60-65 °F (15-18 °C)

### Medien

Use a well-drained, growing substrate with 10-30 % clay, 0-15 % parts (e.g. bark, sand, perlite), 1-2 kg /m<sup>3</sup>, complete balanced fertilizer. 2-4 kg/m<sup>3</sup> slow release fertilizer (3-9 months), iron-chelate, micronutrients, pH 5.5-6.5. Field: well-drained, moderate nutrition supply, humus, loamy soils. Standard fertilization: 60-80 g/m<sup>2</sup> slow release fertilizer (potassium balanced). Do not improve the soil with peat. Keep the soil open.

### Temperatur

Grow at 15-18 °C (59-64,5 °F) or outdoors. In winter indoors frost free at 3-5 °C (37,5-41 °F) or outdoors. Outdoor fleece cover and dry substrate needed. In spring the plants start to grow for 10-12 weeks at 15-18 °C (59-64,5 °F). A chilling period for flower initiation is not required.

### Düngung

High fertilization levels are required. Fertilize the crop weekly with 100-150 ppm nitrogen (at 2 kg/m<sup>3</sup> slow release fertilizer in substrate), using a complete balanced fertilizer. Avoid high ammonium and high nitrogen levels. Don't fertilize after flowering or after early September. In spring fertilize with 100-150 ppm nitrogen using a complete balanced fertilizer. Prevent magnesium deficiency by applying magnesium sulphate (0,05 %) 1-2 times and in case of Iron deficiency apply iron-chelate for 1-2 times. The fertilization with slow release fertilizer is preferred to fertilization with complete balanced fertilizer, because the roots are very sensitive to high salt levels in substrates. Avoid high fertilizer concentrations, it is advisable to fertilize several times with low concentrations weekly. Field: After harvest fertilize the plant additionally with 60-80 g/m<sup>2</sup> of a slow release and potassium balanced fertilizer. Don't fertilize after flowering.

---

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

## FARBEN DER SERIE

Delphinium hybrida Benarys Pacific



**Astolat**  
DH0109R



**Black Knight**  
DH0101R



**Blue Bird**  
DH0106R



**Blue Jay**  
DH0107R



**Cameliard**  
DH0102R



**Galahad**  
DH0110R



**Guinevere**  
DH0108R



**King Arthur**  
DH0104R



**Percival**  
DH0105R



**Summer Skies**  
DH0103R



**Mix**  
DH0199R