

Platycodon grandiflorus F₁

Pop Star

White

Art.-Nr.: PG0103R



A Pop Star is born!

- FastraX perennial: First year flowering without vernalization
- Most compact and well-branched habit on market
- Tight flowering window
- Low requirement of PGR application – less chemicals, more sustainable, cost and time saving
- Use as indoor and outdoor plant

Technical Guide: [Click here](#)

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

Crop Time	Frühjahr: 13 - 15 Wochen
Höhe 	18 cm
Breite 	18 cm
Standort	Sun - Partial shade
Seed Form	Roh
Heat Zone	9-1
Hardiness Zone	3b-9b
Best Uses	Beet und Balkon, Landschaft

KULTURANLEITUNG

Platycodon grandiflorus F₁ Pop Star

Verwendung

Pots, Mixed Containers and Landscape

Aussaattermin

January – May

Keimbedingung

Optimal conditions for seedling development, beginning on the day of sowing until radical emergence. Expect radicle emergence in 3-4 days.

Medien

Plug Culture:

Begin with a Saturated (5) media for the first 4 days. On day 5 begin to reduce the moisture level to a Wet (4) for the next 4 - 5 days. Once the cotyledons have expanded reduce further to a Moist (3). This should occur on day 11-12. Begin to alternate between a moisture level Wet (4) and a Medium (2). Let the media approach a Medium (2) before re-saturating to a Wet (4). Alternate between moisture levels Wet (4) and Medium (2). Allow the media to approach a Medium (2) before re-saturating to a Wet (4). Platycodon prefer slightly drier media conditions for good root development.

Growing On:

Alternate between moisture levels Wet (4) and Medium (2). Allow the media to reach a Medium (2) before re-saturating to a Wet (4). Allowing the media moisture level to dry back will encourage good root development.

Temperatur

Plug Culture:

68-70 °F (20-21 °C), after germination has occurred the temperature can be reduced slightly to 65-68 °F (18- 20 °C).

Growing On:

65-68 °F (18-20 °C) nights, 70-74 °F (21-23 °C) days for the first two weeks after transplanting. Thereafter temperatures may be lowered to 62-65 °F (16-18°C) day and night. An ADT (average daily temperature of 67 °F (19 °C) will give the fastest finished crop.

Düngung

Plug Culture:

Maintain an EC < 1.0. Fertilized water should not exceed an EC of 0.5. Begin fertilizing early using a calcium-based feed, 14-4-14 or 15-5-15 at 50 – 60 ppm.

Growing On:

Higher rates of ammonium can now be used in the feed program. Fertilize at 150 - 200 ppm N using a 17-5-17 or 20-10-20 fertilizer. Under higher light conditions 20-10-20 can be used.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

FARBEN DER SERIE

Platycodon grandiflorus F₁ Pop Star



Blue
PG0101R



Pink
PG0102R



White
PG0103R