

Viola wittrockiana F₁



Art.-Nr.: VW0704R/E



The Mounding Must-Have for Spring and Fall!

- Mounding, globe shaped plant habit
- A multi-purpose, not day length sensitive Pansy
- Enormous, long-lasting flower power
- Suitable for premium pots and baskets

Technical Guide: Click here

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

Crop Time	Frühjahr: 23 - 25 Wochen , Herbst: 11 - 13 Wochen
Höhe ø	13 cm
Breite Ø	18 cm
Standort	Sun
Seed Form	Roh, BeGreen Priming
Best Uses	Ampel



KULTURANLEITUNG

Viola wittrockiana F₁ Highflyer™

Verwendung

Baskets

Kornablage

1 seed per plug. Cover lightly with a thin layer of coarse vermiculite.

Topfen

Viola overall is a day length neutral plant that will flower regardless of day length. Highflyer[™] also flowers under short day conditions below 8 hours, allowing for later sowings and a continuous flowering all winter and spring long.

Medien

Plug culture: pH 5.5-5.8; EC 0.5. Begin with saturated (5) for days 1-5 and then reduce to a moist (3) on day 6. As the seedlings become fully developed with expanded cotyledons the moisture level can be decreased further to a medium (2) on day 9. At this point alternate between a wet (4) and a medium (2) watering.

Growing on: pH 5.5-5.8; keep the pH in the lower range; EC 1.25-1.5. Alternate between moisture levels wet (4) and medium (2).

Temperatur

18-20 °C until root emergence, then lower the temperature gradually to 17-18 °C. After transplanting, maintain 18 °C nights, 18-21 °C days. When seedlings are well established the night temperature can be lowered to 15 °C to tone the plants as flower initiation occurs.

Düngung

Plug culture: Begin feeding early using a calcium based fertilizer at lower rates to keep an adequate amount of calcium and nitrogen supplied to the seedlings. On days 5-7 begin feeding with a calcium-based fertilizer (14-2-14; 13-2-13; 15-5-15 or 17-5-17) at 50-60 ppm. Maintain the EC between 0.5 and 0.75.

Growing on: Fertilize with a calcium-based feed 14-4-14, 15-5-15 or 17-5-15 at 100-150 ppm as needed. Keeping the EC below 1.5 will help prevent root problems.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons. Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair. Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable. Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.



FARBEN DER SERIE

Blue Heaven VW0707R/E



Yellow Purple Wing VW0704R/E



Lemon Splash VW0706R/E



Yellow Red Wing IMPROVED VW0703R/E



Light Violet Shades VW0705R/E



Mix VW0799R/E



Viola wittrockiana F₁ Highflyer™

Violet Face VW0708R/E



White VW0701R/E



Yellow VW0702R/E