

Aquilegia hybrida F₁

Spring Magic®

Rose & Ivory

Art.-Nr.: AH0106R



- Early spring flowering perennial
- Perfect combination with Pansy production
- Dense leaf rosettes in full color range
- Full, globe shaped plant habit
- Low heat requirement
- BeGreen Primed: Chemical and microplastic-free

Crop Time	Frühjahr: 22 - 26 Wochen
Höhe 	35 cm
Standort	Sun
Seed Form	Roh
Heat Zone	9-3
Hardiness Zone	3-8
Best Uses	Beet und Balkon, Steingarten

KULTURANLEITUNG

Aquilegia hybrida F₁ Spring Magic®

Verwendung

Perennial beds and borders, spring bedding, seasonal mixed containers

Aussaattermin

Indoor forcing: Mid May-End July for flowering pots the following year (February onwards); Outdoor forcing: March-April for flowering pots the following year (April onwards)

Kornablage

2-3 seeds per plug

Keimbedingung

14-21 days at 72-77 °F (22-25 °C), in media with low soluble salt levels below 0.5 and a pH of 5.8-6.2 Maintain uniformly moist soils and humidity levels above 95%. Lights aid in germination.

Topfen

Transplant into a well drained media with a pH of 5.5 to 6.2. Two weeks after transplant begin feeding at 150 to 200 ppm N in well balanced fertilizer mix. Keep EC levels below 1.0. Flowers initiate when mature plants are grown below 50°F (10°C) or plugs are treated with temperatures below 40°F (5°C). Vernalization requirements vary by variety and plant maturity. Plant growth is optimum between 60-68°F (15-18°C) nights. Temperatures below 55°F (13°C) slow plant development and extend bloom time and intensify flower colour.

Medien

Use a well-drained, perennial substrate with 0-15 % clay, 0-15 % organic parts (e.g. wood fibres, bark), 1-1,5 kg/m³, complete balanced fertilizer, 1-2 kg/m³ slow release fertilizer (3-9 months), iron-chelate, micronutrients, pH: 5.5-6.2.

Temperatur

Grow at 14-16°C. After the roots are developed decrease the temperature to 8-10°C. The temperature change will support the plant quality. In winter indoors frost free at 3-5°C or outdoors. Forcing in spring for 6-7 weeks at 15-18°C. Cold temperature at 10-12°C will increase the cultivation time. Avoid high temperature at low light levels. A. hybrida needs for vernalization a chilling period (outdoors or chamber) of at least 6-8 weeks at 3-5°C. The foliage of the plants has to be well developed for the chilling period.

Düngung

Moderate fertilization levels are required. Fertilize the crop weekly with 130 -150 ppm nitrogen (at 2 kg/m³ slow release fertilizer in substrate), using a potassium balanced fertilizer (N: K₂O-ratio: 1:1,5). Avoid high ammonium and high nitrogen levels. Don't fertilize after mid September. In spring fertilize with 100-150 ppm nitrogen of a potassium balanced fertilizer.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

FARBEN DER SERIE

Aquilegia hybrida F₁ Spring Magic®



Blue & White
AH0101R



Navy & White
AH0102R



Pink & White
AH0104R



Rose & Ivory
AH0106R



Rose & White
AH0105R



White
AH0107R



Yellow
AH0103R



Mix
AH0199R