



Begonia semperflorens F<sub>1</sub>

# Super Olympia®

#### **Rose**

Art.-Nr.: BS0304P



- Vigorous semperflorens begonia
- Suitable for landscape application
- Great field performance
- Performs in widest array of conditions
- Green foliage partner to Cocktail<sup>®</sup>

#### **Technical Guide: Click here**

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

Crop Time	Frühjahr: 12 - 14 Wochen
Höhe Ø	28 cm
Breite Ø	23 cm
Standort	Sun - Shade
Seed Form	Pilliert
Best Uses	Beet und Balkon, Landschaft



## **KULTURANLEITUNG**

Begonia semperflorens F<sub>1</sub> Super Olympia®

#### Verwendung

Annual, packs, pots, containers and landscape

#### Aussaattermin

December onwards

#### Kornablage

1-2 seeds/pills per plug

#### Keimbedingung

Light is required for germination. 5-10 days at temperatures of 75-80 °F (24-26 °C), and relative humidity levels of 95% or higher. For pelleted seed, slightly higher temperatures and 100% relative humidity aid in uniformity of germination. pH levels of 5.5-6.5 are optimum.

#### **Topfen**

Transplant plugs after 7-9 weeks and grow on at 62-68 °F (16-20 °C). Fertilize weekly with 150 ppm nitrogen watching to avoid excessive salt buildups.

#### Medien

Use a well-drained, growing substrate, pH 5.5-6.2.

#### Temperatur

Grow at 17-19 °C. Before selling harden plants at 7-10 °C and get them used to UV light. Begonia semperflorens does not tolerate frost.

#### Düngung

Moderate fertilization levels are required. Fertilize weekly with 100-150 ppm nitrogen, using a complete balanced fertilizer. High ammonium and high nitrogen levels cause big foliage. Avoid pH above 6.5, as high pH causes iron deficiency. Apply Fe-chelate if chlorosis becomes a problem. Avoid high salt levels in substrate.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

 $Stage\ II\ Starts\ from\ fully\ developed\ cotyledons.\ Ends\ with\ the\ fully\ developed\ true\ leaf\ pair.$ 

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.



# **FARBEN DER SERIE**

## Begonia semperflorens F<sub>1</sub> Super Olympia®







Pink BS0302P



Red BS0303P



Rose BS0304P



White BS0305P



Mix BS0399P