

Lobelia erinus F₁

Masterpiece

Masterpiece — The First Lobelia F₁ Hybrid from Seed!



- Excellent heat tolerance and impressive field performance
- Early flowering with large flowers
- Well-branched, mounding growth
- Perfect for pot and container production

Multipellets only available in North America.

Technical Guide: [Click here](#)

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

Crop Time	Frühjahr: 12 - 14 Wochen
Höhe 	30 cm
Breite 	40 cm
Standort	Sun - Partial shade
Seed Form	Pilliert
Best Uses	Landschaft, Topfpflanze

KULTURANLEITUNG

Lobelia erinus F₁ Masterpiece

Verwendung

Pots, containers, baskets and landscape

Aussaattermin

From January onwards

Kornablage

4 pellets per plug. Light is required for germination - no covering.

Keimbedingung

Optimum conditions for seedling development, beginning on the day of sowing until radicle emergence. Expect radicle emergence in 4-7 days. Humidity should be between 95-100% until day 11, then reduce to 40-60%.

Topfen

Facultative long day plant. Flowering is affected by day-length, irradiance and temperature. Under long day length above 12 hours, the plants will initiate and flower more quickly.

Medien

Plug culture: pH 5.8-6.2; EC 0.5-0.75. Begin with a wet (4) for the first 10 days and on day 11 begin to dry them back slightly to moist (3).

Growing on: pH 5.8-6.2; EC 1.25-1.5. Alternate between moisture levels moist (3) and medium (2). Let plants dry back to at least a medium (2) before re-saturating to a moist (3). Extremely dry plants will have a grayish cast to the leaves. Avoid watering plants under high temperature and light when the leaf temperature is excessive. This can cause leaf burn. Providing good ventilation and horizontal airflow will help lower the humidity and dry back the media, providing oxygen to the roots.

Temperatur

Plug culture: 20-22 °C for days 1-11. For irrigation use warm water above 15 °C only. After development of the roots, the temperature can be further decreased to 18-20 °C.

Growing on: After transplanting, grow at night temperatures of 13-16 °C and day temperatures of 19-22 °C. Lower temperatures may significantly reduce the growth of the plants, but result in a higher plant quality.

Düngung

Start the first fertilization approximately two weeks after sowing to prevent deficiency symptoms. Maintain an EC < 0.7 and begin feeding with less than 100 ppm nitrogen. Increase fertilization in the second part of the plug time up to 100-175 ppm nitrogen. Feed the plants weekly with 175-225 ppm nitrogen, using a complete balanced fertilizer such as 15-5-15 or 18-11-18. Maintain an EC between 1.2 and 1.5. Avoid high ammonium and high nitrogen levels. Wetness and cold medium temperatures are a cause for iron deficiency. The roots are sensitive to high salt levels in substrate.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

FARBEN DER SERIE

Lobelia erinus F₁ Masterpiece



Blue with Eye
LE0201P