


Craspedia globosa

SolarPop

Art.-Nr.: CG0201R



- Large, golden flower heads
- Trendy highlight in every flower bouquet
- Attracts many beneficial insects
- Suitable for pots, containers and the landscape

Crop Time	Frühjahr: 12 - 14 Wochen
Höhe 	65 cm
Breite 	35 cm
Standort	Sun - Partial shade
Seed Form	Roh
Product Use	Topfpflanze, Containers, Landschaft, Schnittblume
Family, Origin	Asteraceae, Australia & New Zealand
Minimum Germ. Rate	85%

TECHNICAL GUIDE

Craspedia globosa SolarPop

Verwendung

Pots, Containers, Cut Flower and Landscape

Kornablage

3 seeds per plug.

Keimbedingung

Optimum conditions for seedling development, beginning on the day of sowing until radicle emergence. Expect radicle emergence in 4-7 days.

Temperatur

Plug culture: 20-22 °C (68-72 °F) on days 1-7. For irrigation use warm water (above 18 °C/ 64 °F) only. 18-20 °C (64-68 °F) during night and day. When the roots reach the bottom of the cell, the temperature can be lowered to 14-16 °C (57-60 °F). Cooler temperatures promote uniformity and compactness of the plugs.

Growing on: 14-16 °C (57-60 °F) days, 12-14 °C (54-57 °F) nights for two weeks after transplanting. After that, temperatures can be reduced by 2 °C (36 °F) day and night.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

FARBEN DER SERIE

Craspedia globosa SolarPop



CG0201R